

FILE COPY

Roy W. Simonson, Director, Soil Classification  
& Correlation, SCS, Hyattsville, Maryland

July 28, 1967

Andrew R. Aandahl, Principal Soil Correlator,  
Midwest Region, SCS, Lincoln, Nebraska

SOILS - Classification and correlation of the soils of Howard County,  
Indiana

Attached are two copies of the classification and correlation of the  
soils of Howard County, Indiana. Copies have been distributed as  
follows:

R. W. Simonson	2 copies
T. B. Evans	6 copies
H. P. Ulrich	2 copies
B. A. Barnes	5 copies

The following revised drafts of descriptions are being sent to you  
along with the final correlation:

Blount	Genesee	Pewamo
Brookston	Kokomo	Russell
Crosby	Miami	
Fincastle	Ockley	

One copy of the soils handbook is also enclosed. Also one set of  
correlation cards are attached.

Attachments

cc:

C. D. Jaedicke  
T. B. Evans (6)  
H. P. Ulrich (2)  
B. A. Barnes (5)  
A. J. Baur (1)  
L. J. Bartelli (1)  
W. M. Johnson (1)  
w/attachments

UNITED STATES DEPARTMENT OF AGRICULTURE  
Soil Conservation Service  
Midwest Regional Technical Service Center  
Lincoln, Nebraska 68508

Classification and Correlation

of the

Soils of Howard County, Indiana

This correlation was prepared in conference by H. Ulrich, F. Sanders, J. Deal, K. Huffman and R. I. Turner during the week of March 20-24, 1967. Evidence used consisted of the soils handbook, revised series descriptions, field correlation, correlation samples, and limited laboratory data.

<u>Symbols</u>	<u>Approved Name</u>
732-A-1	Blount silt loam, 0 to 2 percent slopes
732-A-0	
732-B-1	Blount silt loam, 2 to 4 percent slopes, eroded
732-B-2	
5148-A-0	Brookston silty clay loam
5148-B-0	
148-A-0	
3148-A-+	
3148-A-0	
148-A-+	
140-A-0	Carlisle muck
140-A-+	
300-A-0	
5142-A-1	Crosby silt loam, 0 to 2 percent slopes
142-A-1	
142-A-2	
142-A-+	
142d-A-1	
182-A-1	
142-A-0	
142-B-1	Crosby silt loam, 2 to 4 percent slopes, eroded
142-B-2	
143-B-1	Crosby-Miami silt loams, 2 to 6 percent slopes, eroded
143-B-2	

Symbols	Approved Name
372-A-+	Fincastle silt loam
372-A-1	
372-A-2	
482-A-1	
442-A-1	
372-B-2	
335-A-1	Fox silt loam, 0 to 2 percent slopes
5335-A-1	
145-A-1	
335-B-2	Fox silt loam, 2 to 6 percent slopes, eroded
5335-B-2	
145-B-2	
335-B-3	
335-B-1	
335-C-1	Fox soils, 6 to 12 percent slopes, severely eroded
335-C-2	
335-C-3	
5335-C-3	
5145-C-3	
145-C-2	
145-C-3	
335-D-3	
5335-D-3	
145-D-2	
335-D-2	
445-C-3	
5014-A-0	Genesee silt loam
14-A-0	
6014-A-0	
13-A-0	
Gravel pits	Gravel pits
GP	
146-F-1	9336-F-1 Hennepin loam, 25 to 60 percent slopes
736-F-1	9336-G-1
146-F-3	336-F-1
146-G-1	336-E-1
736-G-1	336-E-2
734-E-3	335-E-1
144-E-1	335-E-3
146-E-1	144-E-3
144-E-2	146-E-3
146-E-2	

Symbols	Approved Name
3149-A-0	Kokomo silty clay loam
3149-A-+	
3489-A-0	
3489V-A-0	
108-A-0	Kokomo silt loam, overwash
5140-A-0	Linwood muck
5140-A-+	
7140-A-0	
1	Made land
144-B-1	Miami silt loam, 2 to 6 percent slopes, eroded
144-B-2	
5144-B-2	
144-B-0	
144-C-1	Miami silt loam, 6 to 12 percent slopes, eroded
144-C-2	
143-C-2	
144-C-0	
184-C-2	
144-D-1	
144-D-2	
144-D-0	
374-C-1	
144-B-3	Miami clay loam, 2 to 6 percent slopes, severely eroded
184-E-3	
143-B-3	
144-C-3	Miami clay loam, 6 to 12 percent slopes, severely eroded
184-C-3	
144-D-3	Miami clay loam, 12 to 18 percent slopes, severely eroded
184-D-3	
734-B-1	Morley silt loam, 2 to 6 percent slopes, eroded
734-B-2	
734-B-3	Morley silty clay loam, 2 to 6 percent slopes, severely eroded
734-C-1	Morley silty clay loam, 6 to 12 percent slopes,
734-C-2	severely eroded
734-C-3	
734-D-2	
734-D-3	

Symbols	Approved Name
445-A-1	Ockley silt loam, 0 to 2 percent slopes
445-A-+	
445-B-1	Ockley silt loam, 2 to 6 percent slopes, eroded
445-B-2	
445-C-1	
445-C-2	
184-A-1	Ockley silt loam, loamy substratum, 0 to 2 percent slopes
184-B-1	Ockley silt loam, loamy substratum,
184-B-2	2 to 6 percent slopes, eroded
3488V-A-0	Patton silty clay loam, loamy substratum
3488-A-0	
3448-A-0	
3018-A-0	Patton silty clay loam, occasionally flooded
3738-A-+	Pewamo silty clay loam
3738-A-0	
3739-A-0	
Quarries ↗	Quarries
Stone quarries	
Limestone quarries	
374-A-1	Russell silt loam, 0 to 2 percent slopes
373-A-1	
484-A-1	
144-A-1	
143-A-1	
374-B-1	Russell silt loam, 2 to 6 percent slopes,
374-B-2	moderate erosion eroded
484-B-2	
374-C-2	
373-B-1	
373-B-2	
12-A-0	Shoals silt loam
18-A-0	

Approved: 7-31-67

*AR Aandahl*  
 Andrew R. Aandahl, Principal Soil  
 Correlator, Midwest Region

## CLASSIFICATION OF SERIES

Howard County, Indiana

Blount	Aeric Ochraqualfs; fine, illitic, mesic.
Brookston	Typic Argiaquolls; fine-loamy, mixed, noncalcareous, mesic.
Carlisle	Organic.
Crosby	Aeric Ochraqualfs; fine-loamy, mixed, mesic.
Fincastle	Aeric Ochraqualfs; fine-silty, mixed, mesic. (Typic Ochraqualfs?)
Fox	Typic Hapludalfs; fine-loamy over sandy or sandy-skeletal, mixed, mesic.
Genesee	Fluventic Eutrochrepts; fine-loamy, mixed, mesic.
Hennepin	Typic Eutrochrepts; fine-loamy, mixed, mesic.
Kokomo	Typic Argiaquolls; fine, mixed, noncalcareous, mesic.
Linwood	Organic.
Miami	Typic Hapludalfs; fine-loamy, mixed, mesic.
Morley	Typic Hapludalfs; fine, illitic, mesic.
Ockley	Typic Hapludalfs; fine-loamy, mixed, mesic.
Patton	Typic Haplaquolls; fine-silty, mixed noncalcareous, mesic.
Pewamo	Typic Argiaquolls; fine, mixed, noncalcareous, mesic.
Russell	Typic Hapludalfs; fine-silty, mixed, mesic.
Shoals	Aeric Fluventic Haplaquepts; fine-loamy, mixed, nonacid, mesic. (Fluventic Haplaquepts?)